



Prescribing Guidelines for Urinary Tract Infections

PARTNERS
FOR KIDS®





Urinary Tract Infections in Children

Urinary tract infections (UTIs) are common infections of childhood that may affect any part of the urinary tract, from the urethra to the kidneys. The following guidance will focus on treating uncomplicated UTIs with an appropriate narrow spectrum antibiotic for the shortest effective duration. Infections in children with underlying renal diseases, anatomic or functional anomalies of the urinary tract, urinary catheters, or immunocompromising conditions/therapies constitute complicated UTIs and must be approached on an individual basis.


General Management and Treatment

1. Start by obtaining an appropriate urine sample for urinalysis and urine culture. Urine culture should be performed if urinalysis is positive or if there is a high clinical suspicion for infection.
 - a. Clean-catch urine specimens should be obtained in all toilet trained children. While bagged urine samples can be used for initial urinalysis, catheterization is recommended to obtain appropriate samples for urine cultures in pre-toilet trained children.
2. Decide whether empiric antibiotic therapy for presumed UTI is warranted, and prescribe the most appropriate empiric therapy for the optimal duration (see Table 1).
 - a. Does the child have presumed cystitis or pyelonephritis?
 - i. These symptoms suggest pyelonephritis and it is recommended to treat empirically if present: Fever, vomiting, flank pain, or costovertebral angle tenderness.
 - ii. Children without signs or symptoms of pyelonephritis: Consider waiting for urine culture results to make a definitive diagnosis and optimal management.
 - b. Has the child had prior culture-confirmed UTIs to guide empiric therapy?
 - i. Previous urine culture results and susceptibilities may help guide the choice of empiric therapy.

Table 1: Empiric Antibiotic Therapy

Suspected Diagnosis	Age Range	Preferred Antibiotic*
Bacterial Cystitis	<12 years	Cephalexin 25 mg/kg/dose BID x 3 days (max 500 mg/dose)
	≥12 years	Nitrofurantoin macrocrystal/monohydrate 100 mg BID x 3 days ^c
Pyelonephritis	<12 years	Cephalexin 25 mg/kg/dose TID x 7 days (max 1000 mg/dose) ^b
	≥12 years	Cephalexin 1000 mg TID x 7 days ^b
<p>a. Preferred empiric antibiotic choices are based on commonly recovered pathogens, antimicrobial susceptibility patterns at Nationwide Children's Hospital, and drug penetration into sites of infection. It is recommended to review local resistance patterns and antibiograms for susceptibility in your area</p> <p>b. See cefdinir dosing in 'Culture Directed Therapy' if cephalexin is not an option due to adherence concerns</p> <p>c. Cephalexin 500 mg BID x 3 days is an alternative therapy if nitrofurantoin is not an option</p>		

Table 2: Culture-directed Therapy

Spectrum	Antibiotic	Dose	Frequency and Duration	
			Cystitis	Pyelonephritis
Most Preferred 	Nitrofurantoin ^a	100 mg/dose (adult dose)	BID x 3 days	Do NOT Use
	Amoxicillin	20 mg/kg/dose (max 875 mg/dose)	BID x 3 days	TID x 7 days
	Cephalexin	25 mg/kg/dose	BID x 3 days (max 500 mg/dose)	TID x 7 days (max 1000 mg/dose)
	TMP/SMX	4 mg/kg/dose (max 160 mg/dose)	BID x 3 days	BID x 7 days
	Cefdinir	7 mg/kg/dose (max 300 mg/dose)	BID x 3 days	BID x 7 days
	Amoxicillin/ Clavulanate ^b	20 mg/kg/dose (max 875 mg/dose) ^b	BID x 3 days	TID x 7 days
	Ciprofloxacin	15 mg/kg/dose (max 750 mg/dose)	BID x 3 days	BID x 7 days
Least Preferred				
a. Macrocrystal/Monohydrate (age ≥12 years). Use for cystitis only; Nitrofurantoin should NOT be used for Pyelonephritis/Febrile UTI b. Dose based on amoxicillin component. Use products with 7:1 ratio of amoxicillin to clavulanate below: Suspension: 400 mg/57 mg/5 mL Tablet: 875 mg/125 mg Chewable tablet: 400 mg/57 mg				

3. Follow-up of urine culture results

- For follow-up management, adjust the antibiotic therapy as indicated depending on the results of the urine culture and susceptibilities (see Table 2). If the urine culture does not suggest a UTI, stop the antibiotic and consider other diagnoses.
- For children who improve with treatment, test-of-cure urine cultures are NOT routinely recommended.

4. For children who do not improve within 48-72 hours of appropriate antibiotic therapy, consider obtaining a renal/bladder ultrasound to evaluate for complications such as renal or perirenal abscess.

5. When to refer:

- Consider referral to Pediatric Urology for further evaluation of children with recurrent UTIs, abnormal renal/bladder ultrasound results, or symptoms of voiding dysfunction when not infected (urgency, frequency, enuresis).



References

1. Roberts KB, Subcommittee on Urinary Tract Infection, Steering Committee on Quality Improvement and Management. Urinary Tract Infection: Clinical Practice Guideline for the Diagnosis and Management of the Initial UTI in Febrile Infants and Children 2 to 24 Months. *Pediatrics*. 2011;128(3):595-610.
2. Shaikh N, Morone NE, Lopez J, et al. Does this child have a urinary tract infection? *JAMA*. 2007;298(24):2895-2904.
3. Montini G, Tullus K, Hewitt I. Febrile urinary tract infections in children. *N Engl J Med*. 2011;365(3):239-250.
4. Shaikh N, Mattoo TK, Keren R, et al. Early Antibiotic Treatment for Pediatric Febrile Urinary Tract Infection and Renal Scarring. *JAMA Pediatr*. 2016;170(9):848-854.
5. Edlin RS, Shapiro DJ, Hersh AL, Copp HL. Antibiotic resistance patterns of outpatient pediatric urinary tract infections. *J Urol*. 2013;190(1):222-227.
6. Keren R, Chan E. A meta-analysis of randomized, controlled trials comparing short- and long-course antibiotic therapy for urinary tract infections in children. *Pediatrics*. 2002;109(5):E70-70.
7. Michael M, Hodson EM, Craig JC, Martin S, Moyer VA. Short compared with standard duration of antibiotic treatment for urinary tract infection: a systematic review of randomised controlled trials. *Arch Dis Child*. 2002;87(2):118-123.
8. Shaikh N, Hoberman A. Urinary tract infections in infants older than one month and young children: Acute management, imaging, and prognosis. In: Torchia MM, ed. UpToDate. Waltham, MA: UpToDate.
9. Saadeh SA, Mattoo TK. Managing urinary tract infections. *Pediatr Nephrol*. 2011;26(11):1967-1976.
10. National Collaborating Centre for Women's and Children's Health. *Urinary tract infection in children: diagnosis, treatment and long-term management*. London, UK: National Institute for Health and Clinical Excellence;2007.
11. Stein R, Dogan HS, Hoebeke P, et al. Urinary tract infections in children: EAU/ESPU guidelines. *Eur Urol*. 2015;67(3):546-558.
12. Working Group of the Clinical Practice Guidelines for Urinary Tract Infection in Children. *Clinical Practice Guidelines for Urinary Tract Infection in Children. Ministry of Health National Health Service Quality Plan, Social and Equality Policy; Aragon Institute of Health Sciences (I+CS) 2011, SNS Clinical Practice Guidelines: I+CS No 2009/01*.
13. Hoberman A, Wald ER. Urinary tract infections in young febrile children. *Pediatr Infect Dis J*. 1997;16(1):11-17.
14. Hoberman A, Greenfield SP, Mattoo TK, et al. Antimicrobial prophylaxis for children with vesicoureteral reflux. *N Engl J Med*. 2014;370(25):2367-2376.
15. Reaffirmation of AAP Clinical Practice Guideline: The Diagnosis and Management of the Initial Urinary Tract Infection in Febrile Infants and Young Children 2-24 Months of Age. *Pediatrics*. 2016;138(6).
16. Shaikh N, Hoberman A, Keren R, et al. Recurrent Urinary Tract Infections in Children With Bladder and Bowel Dysfunction. *Pediatrics*. 2016;137(1).
17. Zaoutis T, Shaikh N, Fisher BT, et al. Short-Course Therapy for Urinary Tract Infections in Children: The SCOUT Randomized Clinical Trial. *JAMA Pediatr*. 2023;177(8):782-789.

18. National Institute for Health and Care Excellence. (2018). Urinary tract infection (lower): antimicrobial prescribing [NG109]. <https://www.nice.org.uk/guidance/ng109>
19. Working Group of the Clinical Practice Guidelines for Urinary Tract Infection in Children. Clinical Practice Guidelines for Urinary Tract Infection in Children. Ministry of Health National Health Service Quality Plan, Social and Equality Policy; Aragon Institute of Health Sciences (I+CS) 2011, SNS Clinical Practice Guidelines: I+CS No 2009/01.
20. Fox MT, Amoah J, Hsu AJ, Herzke CA, Gerber JS, Tamma PD. Comparative Effectiveness of Antibiotic Treatment Duration in Children With Pyelonephritis. *JAMA Netw Open*. 2020 May 1;3(5):e203951.
21. Yu D, Bio LL. Shedding Light on Amoxicillin, Amoxicillin-clavulanate, and Cephalexin Dosing in Children from a Pharmacist's Perspective. *J Pediatric Infect Dis Soc*. 2022 Dec 28;11(12):594-602.
22. Lexicomp Online, Pediatric and Neonatal Lexi-Drugs Online. Waltham, MA: UpToDate, Inc.; November 15, 2023. <https://online.lexi.com>.



Referrals and Consultations

Online: [NationwideChildrens.org/Urology](https://www.NationwideChildrens.org/Urology)

Phone: (614) 722-6200 or (877) 722-6220 | Fax: (614) 722-4000

Physician Direct Connect Line for 24-hour urgent physician consultations:
(614) 355-0221 or (877) 355-0221



PARTNERS
FOR KIDS®

